



National
Library
of Medicine

PubMed	Nucleotide	Protein	Genome	Structure	PMC	Taxonomy	OMIM
Search PubMed	<input type="text"/> for <input type="button" value="Go"/>						
	Limits	Preview/Index	History	Clipboard	Details		
	<input type="button" value="Display"/>	<input type="button" value="Abstract"/>	<input type="button" value="▼ Show: 20"/>	<input type="button" value="▼ Sort"/>	<input type="button" value="▼ Send to: Text"/>	<input type="button" value="▼"/>	

Entrez PubMed

1: J Biomol NMR. 2003 Nov;27(3):193-203.

[Related Articles](#), [Links](#)

1H(C) and (1)H(N) total NOE correlations in a single 3D NMR experiment. (15)N and (13)C time-sharing in t(1) and t(2) dimensions for simultaneous data acquisition.

PubMed Services

Xia Y, Yee A, Arrowsmith CH, Gao X.

Department of Chemistry, University of Houston, Houston, TX 77004-5003, U.S.A. Ontario Cancer Institute and Department of Medical Biophysics, The University of Toronto, Toronto, ON, Canada M5G 2M9.

Simultaneous data acquisition in time-sharing (TS) multi-dimensional NMR experiments has been shown an effective means to reduce experimental time, and thus to accelerate structure determination of proteins. This has been accomplished by spin evolution time-sharing of the X and Y heteronuclei, such as (15)N and (13)C, in one of the time dimensions. In this work, we report a new 3D TS experiment, which allows simultaneous (13)C and (15)N spin labeling coherence in both t(1) and t(2) dimensions to give four NOESY spectra in a single 3D experiment. These spectra represent total NOE correlations between (1)H(N) and (1)H(C) resonances. This strategy of double time-sharing (2TS) results in an overall four-fold reduction in experimental time compared with its conventional counterpart. This 3D 2TS CN-CN-H HSQC-NOESY-HSQC pulse sequence also demonstrates improvements in water suppression, (15)N spectral resolution and sensitivity, which were developed based on 2D TS CN-H HSQC and 3D TS H-CN-H NOESY-HSQC experiments. Combining the 3D TS and the 3D 2TS NOESY experiments, NOE assignment ambiguities and errors are considerably reduced. These results will be useful for rapid protein structure determination to complement the effort of discerning the functions of diverse genomic proteins.

PMID: 12975580 [PubMed - in process]

<input type="button" value="Display"/>	<input type="button" value="Abstract"/>	<input type="button" value="▼ Show: 20"/>	<input type="button" value="▼ Sort"/>	<input type="button" value="▼ Send to: Text"/>	<input type="button" value="▼"/>
--	---	---	---------------------------------------	--	----------------------------------

[Write to the Help Desk](#)

[NCBI](#) | [NLM](#) | [NIH](#)

Department of Health & Human Services
Freedom of Information Act | Disclaimer